



Infrared Spectroscopy and Microorganisms Associated with African Nutmeg (*Monodora myristica*) Seeds Sold in a Municipal Market in Imo State, Nigeria

Eberechukwu M. Maduike^{1*} and Nkiruka C. Anuna²

¹*Department of Animal and Environmental Biology, Faculty of Science, University of Port Harcourt, P.M.B. 5323 Choba, Port Harcourt, Nigeria.*

²*Department of Science Laboratory Technology, Akanu Ibiam Federal Polytechnic, Uwana, Ebonyi State, Nigeria.*

Authors' contributions

This work was carried out in collaboration between both authors. Author EMM designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author EMM managed the analyses of the study. Author NCA managed the literature searches. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRIMPS/2018/40661

Editor(s):

- (1) Edet O. Akpanyung, Professor, Department of Biochemistry, University of Uyo, Nigeria.
(2) Somdet Srichairatanakool , Professor, Department of Biochemistry, Faculty of Medicine, Chiang Mai University, Chiang Mai 50200, Thailand.
(3) BuLang Gao, Professor, Department of Medical Research, Shijiazhuang First Hospital, Hebei Medical University, China and Department of Radiology, Shanghai Jiaotong University Renji Hospital, China.

Reviewers:

- (1) Arunsi Uche Okuu, Abia State University, Nigeria.
(2) Vladimíra Tomečková, University of Pavol Jozef Šafárik in Košice, Slovakia.
Complete Peer review History: <http://www.sciedomain.org/review-history/24445>

Original Research Article

Received 28th January 2018

Accepted 8th April 2018

Published 3rd May 2018

ABSTRACT

This work evaluated the infra-red (IR) spectra of the oil extract from African nutmeg (*Monodora myristica*) seeds, popularly called 'ehuru' in the eastern part of Nigeria – among the Igbo. The microbial analysis of the dried 'ehuru' seeds was also carried out in order to ascertain the prevailing bacteria and fungi on the outer coat of this invaluable seed as sold in the market. *Monodora myristica* oil extract was obtained through soxhlet extractor using ethanol of Analar grade. About 15 g of the extract (oil) was obtained after assessing 400 g of the ground dried seeds

*Corresponding author: E-mail: eberechukwu.maduike@uniport.edu.ng;

of the sample. Infra-red analysis of the oil extract was evaluated. The IR spectra of the extract indicated that the sample contained the following functional groups: phosphate esters, ketones, amides, amines, amino acids, ammonium salts, alkenes, phenols, alkanes, ether, lactams and carboxyl groups. There is no doubt that the presence of these chemical functional groups in *Monodora myristica* seed conferred to its antimicrobial quality, therapeutic potentials, and its use as food additive (spice). Common microorganisms associated with the dried 'ehuru' seeds were bacteria - *Streptococcus* sp. and *Staphylococcus* sp.; fungus - *mucor* sp. The presence of these microorganisms on the seed might have been introduced as a result of exposure to unhygienic conditions by local handlers and frequent touching by buyers.

Keywords: African nutmeg; oil extract; IR spectra; functional group; microorganisms.

1. INTRODUCTION

African nutmeg (*Monodora myristica*) seed, popularly called 'ehuru' among the Igbo ethnic group in Nigeria is one of the commonly used spices in the eastern part of the country. It has the qualities of aromatic seasoning and flavour. It is only added to foods in small amount but it makes important contribution towards the aroma and flavour of foods due to the presence of the volatile oil (essential oil) and fixed oil. The seed is oblong and pale brown when fresh with a thin seed coat and hard kernel [1]. *Monodora myristica* is a flowering perennial edible plant which belongs to the family Annonaceae. It is commonly found in the West African evergreen forests in countries like Nigeria, Cameroon, Liberia, Angola and West Kenya [2,3]. The tree can grow up to a height of about 35 m and 2 m in diameter [4]. The plant starts fruiting between August and November [5]. The fruit *M. Myristica* is a berry of about 20 cm diameter; green and spherical, and becomes woody at maturity. The fruit is attached to a long stalk which is up to 60 cm long. Inside the fruit are the numerous oblong, pale brown seeds which are usually 1.5 cm long and are surrounded by a whitish fragrant pulp [6]. The oil extract from *Monodora myristica* seed contains significant pharmacological compounds like alkaloids, flavonoids, vitamin A and E as well as many vital lipids [7-10]. Traditionally, the plant is widely used especially to relieve toothache as well as in the treatment of dysentery. When roasted and ground, the seeds are rubbed on the skin for the treatment of skin diseases [11]. The seed is used by most Ethnic Nationalities in the various regions of Nigeria, as spice in the preparation of pepper soup (a traditional sauce used for health improvement for the healthy, sick and convalescing). Also, it is reported that natives in the Democratic Republic of Congo use it in the treatment of cough, headache, fever, and

skin diseases. When ground to powder, it is used to prepare pepper soup as stimulant to relieve constipation and control passive uterine haemorrhage in women immediately after child birth. Trado-medically, it has been used to cure mild fever and is known to have diuretic properties [12]. The aromatic stimulating properties of the plant appear to be the reason why it is occasionally used to ease flatulence, vomiting and for the correction of nausea arising from other drugs. *Monodora myristica* has been reported to exhibit antibacterial activity *in vitro* against some human and animal pathogens. This appears to be the bases for the use of the plant in traditional medicine to treat various diseases such as cough, pneumonia, tuberculosis, upper respiratory tract infections fever and skin diseases [12]. The oily extracts of *Monodora myristica* contain phytochemicals such as tannins, glycosides, saponins and Flavonoids which exert antimicrobial effects against *Bacillus subtilis*, *Candida albicans* and *Staphylococcus aureus* [13]. This work was targeted to ascertain the IR spectra of the African nutmeg oil extract and, thus the possible chemical substances present based on functional group. The research also aimed at finding out the possible microbial flora (bacteria and fungi) on the seed coat as sold in the market.

2. MATERIALS AND METHODS

2.1 Collection of Samples

The dried indigenous spice, *Monodora myristica* (ehuru) seeds were purchased from Ekeonunwa market in Owerri, Imo state of Nigeria. Identification and its authentication were done by Odum, D. C. of Department of Science Laboratory Technology (Biology/Microbiology), Federal Polytechnic Nekede, Owerri, Imo State, Nigeria.

2.2 Sample Preparation

The seeds were thoroughly screened to remove the bad ones and stones. The outer seed coats were removed and the good seeds were prepared by grinding them in a laboratory electric mill. The powdered sample was stored in an airtight, sterile glass container and kept in the refrigerator at 4°C until needed for analyses. Some seeds were kept whole (without grinding) for microbial analysis.

2.3 Extraction of Oil

Before extraction, 400 g of the powdered sample of *M. myristica* seed from the refrigerator was heated in a hot-air oven at 60°C in order to remove moisture (for easy extraction of the oil). The powdered seeds were soaked in 1000 ml absolute ethanol (99.9%, boiling range 78.0 - 78.5°C) for 48 hours. The extraction of the oil content of the seed was by soxhlet extractor. After the extraction, the solvent (ethanol) was removed from the extract through the fractional distillation process. At the end of the extraction, 15 g of the extract (oil) was recovered.

2.4 Infrared Spectroscopy

The infrared (IR) spectroscopy was carried out using the Philips Scientific, Model No: 40 13-15-

17494 equipment at University of Uyo Research Laboratory, Akwa-Ibom State, Nigeria.

2.5 Microbial Analysis of the Seed

Nutrient agar (NA) medium was used for the isolation of bacteria on the seeds. This was achieved by aseptically rubbing and placing few of the seeds on the NA agar medium in triplicate plates and was incubated at 30°C for 48 hours. Some seeds were also placed on sabouraud dextrose agar (SDA) in triplicate plates for the isolation of fungi on the seeds. The plates were incubated at 26°C for 4 days. The bacteria isolated were characterized based on cultural characteristics, staining reactions and biochemical reactions as described by Cheesbrough [14], while the morphological characterization of the fungal isolates was through microscopy under lactophenol cotton-blue stain.

3. RESULTS AND DISCUSSION

The infrared (IR) spectroscopy of the oil extract from *M. myristica* was carried out and the spectra analysed following related authors' guidelines [15,16]. Fig. 1 revealed the IR spectra of the oil extract from *M. myristica* seed while Table 1 showed the peak report.




Fig. 1. Infrared spectrum of oil extract from *Monodora myristica* seeds

Table 1. Analysis of infrared spectra of *Monodora myristica* oil extract

Frequency (cm ⁻¹)	Transmittance (%)	Frequency range (cm ⁻¹)	Functional group structure	Inference
1196.91	66.5603	1315 – 1180	p=O	Phosphate ester
1375.96	71.3161	1410 – 1310	O-H	Phenol
1711.36	31.8864	1715 – 1710	C=O	Ketones
3007.96	56.4442	3500 – 3000	N-H	Amines
3141.97	73.4619	3150 – 3050	O-H	Phenol
3266.20	71.8750	3400 – 3200	O-H	Alcohols & Phenols
3345.09	70.8680	3400 – 3200	O-H	Alcohols & Phenols
1244.26	64.2397	1275 – 1200	C-O-C	Ether
1414.67	67.6424	1420 – 1405	C-N	Amides
2854.77	29.7860	3000 – 2850	C-H	Alkanes
3092.48	73.1051	3100 – 2600	N-H	Amino acids
3216.11	72.5560	3300 – 3030	N-H	Ammonium salts
3289.57	71.4642	3300 – 3050	N-H	Amides
3377.59	70.9736	3800 – 3000	N-H	Amines
1279.78	65.8691	1302 – 1034	C-O	Ester
1461.35	58.5447	1650 – 1450	C=C	Alkenes
2925.85	16.1560	3000 – 2500	O-H	Carboxyl
3120.88	73.3944	3150 – 3050	O-H	Phenol
3232.38	72.1722	3300 – 3050	N-H	Amides
3306.22	71.4238	3400 – 3200	N-H	Amides
3392.04	71.2208	3400 – 3200	N-H	Amides & lactams

The microbial flora of *Monodora myristica* seed was studied to investigate the bacterial and fungal diversity associated with the sample as obtainable in the market environment. The result obtained was as shown in Table 2.

Table 2. Microorganisms isolated from the *Monodora myristica* seeds

Bacteria	Fungi
<i>Streptococcus</i> sp.	<i>Mucor</i> sp.
<i>Staphylococcus</i> sp.	

3.1 Discussion

The results obtained from the infrared spectroscopy revealed that *M. myristica* seed contains important organic substances. Phenolic compounds were identified as the most naturally occurring group in the sample. The infrared analysis of *M. myristica* seed by Okwu and Njoku (2009) also showed the presence of phenol (hydroxyl phenol) [17]. These compounds possess a broad spectrum of biochemical activities including antioxidant and radical scavenging properties, and their antioxidant potential is believed to be conferred on them by the presence of a hydroxyl functional group

(-OH), which is bonded directly to an aromatic hydro-carbon (phenyl) ring. This makes them readily donate electrons to free radicals and stabilize them before they destroy living cells [18]. Due to the presence of the phenolic compounds, McCue and Shetty [19] and Etouchi et al. [20] reported that *M. Myristica* extract has the ability to exhibit anti-amylase activities which may be potentially useful in the treatment and management of diabetes mellitus and obesity. Also, the bactericidal and/or bacteriostatic effects of this seed as reported by Odoh et al. [21] and Ogu et al. [22] based on the findings of this study, is as a result of the presence of these phenolic compounds. It has been confirmed that *M. myristica* possesses vermicidal property [9]. This property can be attributed to the presence of phenol and phenolic compounds present in the seed (Table 1).

The study showed that this naturally endowed seed contains ketones (C=O). It has been reported that exogenous ketone supplements may provide benefits which include evident weight loss, athletic performance enhancement, anti-inflammatory properties and cancer prevention [23]. In a study to investigate how african nutmeg (*monodora myristica*) lowers

cholesterol and modulates lipid peroxidation in experimentally induced hypercholesterolemic male wistar rats by Nwozo et al. they reported that the extract was able to control the increase in body weight, thus observed reduction in body weight [6]. It is believed that these aromatic ketones (e.g., gingerol and paradol) give the seed its pungent and peppery taste [24]. Therefore, the African nutmeg seed can be harnessed as an exogenous ketone source for the benefit of mankind directly or indirectly.

Amines (N-H) are very essential in life because they are involved in creating amino acids. Amino acids are the building blocks of proteins in human body. Many vitamins are also made from amino acids. The amine groups which appeared at 3800-3000 cm⁻¹ indicate the presence of alkaloids [17]. It was reported by Okwu (2005) that alkaloids isolated purely from plant extracts and their synthetic derivatives are potential medicinal agents used for fungicidal, bactericidal, antimalarial and analgesic purposes [25]. Amines are mostly used in pharmaceutical industries as analgesics (pain killers), decongestant (agent that relieves nasal congestion), and anesthetics [26]. Thus, the oil extract from this organic seed can be refined and processed for amine production to be used in orthodox medicine.

The IR spectroscopy on the experimental seed revealed that it contains amino acids (N-H). As has been earlier said, amino acids are essential monomers in protein synthesis. It has also been reported that food supplements containing amino acids are good for children's growth, boost the immune system, protect the liver from toxic agents, regulate blood pressure, and accelerate tissue repair [27]. Significant reduction in the levels of serum total cholesterol and triglycerides in hypercholesterolemic rats treated with *M. myristica* oil extract has been reported [6]. This indicates that the extract possesses cholesterol lowering potential. These outlined benefits confirmed its use as therapeutic agent in traditional medicine as reported by Cimanga et al. (2002) [12].

Ether (C-O-C) was another component of the oil extract from African nutmeg. Ether is a highly volatile and flammable liquid that has the ability to depress cerebral activity when used in anaesthetic dosage. Although in some individuals, ether causes irritation to the upper airway, it is a bronchodilator and it can be useful

in treating bronchospasm resistant to other drugs [28]. This confirmed the use of *Monodora myristica* as a curative substance for respiratory tract maladies [12].

Alkane group (C-H) was another observable substance identified in the oil extract of the African nutmeg seed. An investigation study by Olga et al. (2017) involving the utilization of n-alkanes with an odd number of carbon atom as carbon sources by three yeast strain revealed high biomass yield, accumulation of high amount of lipids, and their ability to produce nutritionally important fatty acids [29]. A study with similar alkane group from the oil extract of the seed under investigation may likely produce the same result. Thus, studies to investigate the use of this oil extract to promote and/or achieve desirable traits and characteristics in organisms like fungi and bacteria should be encouraged.

Another functional group indicated in *M. Myristica* seed was the C-O stretching of ester centered at 1302-1034 cm⁻¹. Naturally occurring esters of organic acids in fruits and flowers give them their distinctive smells [30]. Esters with low molecular weight are commonly used as fragrances and found in essential oils. They are also used for food flavourings [31]. Plant esters as contained in *M. myristica* seeds have the ability to lower serum cholesterol by partially inhibiting the absorption of cholesterol in the small intestine [32]. The backbone of DNA molecules are made of phosphate esters hence will be beneficial to human physiological activities.

The reason for the isolation of very few microorganisms from the outer seed coat of *M. myristica* seed is not farfetched (Table 2). This is because an antimicrobial screening done by Adewole et al. [33] using the oil extract from *M. myristica* seed showed that the oil had antimicrobial action against *Escherichia coli*, *Bacillus subtilis* and *Staphylococcus aureus*. It was recorded that the antimicrobial activity exhibited by the oil extract competed favourably with the action of the antibiotic, streptomycin against these bacteria. A similar antimicrobial action on *Staphylococcus aureus*, *Klebsiella pneumonia*, *Escherichia coli* and *Salmonella typhi* was observed using ethanolic extracts of *M. myristica* seed [34]. The identified microorganisms might have been introduced on the seeds of 'ehuru' as a result of exposure to unhygienic conditions by handlers and frequent touching by buyers. It is, therefore, necessary that the seeds be properly

cooked before consumption. There is no doubt that the presence of these chemical functional groups in *M. myristica* seed conferred to it therapeutic, antimicrobial and industrial uses.

4. CONCLUSION

Having confirmed that the oil extract from *Monodora myristica* seed is of great value, large scale production of the seed should be encouraged so that its products like the oil extract can be channelled towards the production of pharmaceutical and personal healthcare products. Pharmaceutical industries are hereby encouraged to harness the great potentials of this nature's gift to mankind. Also, the oil extract from this invaluable seed can be used in the right proportion as dietary supplements to promote healthy living. Therefore, the use of African nutmeg seed oil extract sold as spice, food supplement and health booster should be encouraged. Having seen the various chemical functional groups contained in *Monodora myristica* seed, it is very important that scientists, pharmacist, and dieticians utilize this natural product for antimicrobial, therapeutic, and nutritional purposes.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Okafor JC. Some useful tropical plants in health care delivery. Forestry Commission HQ, Enugu. 2003;3-9.
2. Burabai W, Akor AJ, Igoni AH, Uyate YT. Fracture resistance of African nutmeg (*Monodora myristica*) compressive loading. Electronic Journal of Environmental Agriculture and Food Chemistry. 2007; 6(10):2434-2442.
3. Okafor JC. Development of forest tree crops for food supplies in Nigeria. Forest Ecological Management. 1987;1:235-247.
4. Weiss EA. Spice crops. CABI Publishing, Oxon, UK. 2002;102-103.
ISBN: 13:9780851996059
5. Omobuwajo TO, Omobuwajo OR, Sanni LA. Physical properties of calabash nutmeg (*Monodora myristica*) seeds. Journal of Food Engineering. 2003;57: 375-381.
6. Nwozo SO, Kasumu TF, Oyinloye BE. African nutmeg (*Monodora Myristica*) lowers cholesterol and modulates lipid peroxidation in experimentally induced hypercholesterolemic male wistar rats. International Journal of Biomedical Science. 2015;11(2):86-92.
7. Bassey ME, Johnny II, Okoro BI. Lesser known spices of Akwa Ibom; their nutritional, antinutritional, mineral and phytochemical analysis. Archives of Applied Science Research. 2011;3(3): 553-559.
8. Uhegbu FO, Iweala EJ, Kanu I. Studies on the chemical and anti nutritional content of some Nigerian spices. Inter. J. Nutri. Metab. 2011;3(6):72-76.
9. Ekeanyanwu RC, Etienajirhevwe OF. *In vitro* antihelmintic potentials of *Xylopia aethiopica* and *Monodora myristica* from Nigeria. Afr. J. Biochem. Res. 2012;6(9): 115-120.
10. Nwachukwu N. Nutritional and anti-nutritional substances in some selected indigenous spices. Ph.D Thesis, FUTO, Nigeria; 2000.
11. Irvine FR. Woody Plants of Ghana with special reference to their uses. Oxford University Press, London. 1961;13-23.
12. Cimanga K, Kanbu K, Toma L, Apars S, De Bruyne T, Harnan N, Totte J, et al. Correlation between chemical composition and antibacterial activity of essential oils of some aromatic medicinal plants growing in the Democratic Republic of Congo. Journal of Ethnopharmacology. 2002;79: 213-220.
13. Udoh UE, Ezugwu CO, Ajali LU. Antimicrobial activity of *Monodora myristica* seed oil. Journal of Pharmacology and Allied Sciences. 2004;2(2):233-236.
14. Cheesbrough M. District laboratory practice in tropical countries part 2. Press Syndicate of the University of Cambridge; 2000.
15. Dana WM, Foil AM, Robert WH. Course notes on the interpretation of infrared and

- Raman spectra. John Wiley & Sons, Inc.; 2004.
DOI: 10.1002/0471690082
16. Stuart B. Infrared spectroscopy: Fundamentals and applications. John Wiley & Sons, Ltd.; 2004.
ISBNs: 0-470-85427-8 (HB); 0-470-85428-6 (PB)
17. Okwu DE, Njoku EE. Chemical composition and *in vitro* antifungal activity screening of seed and leaf extracts from *Aframomum melegueta* and *Monodora myristica* against *Sclerotium rolfsii* of cowpea plant (*Vigna unguiculata* L. Walp). Pest Technology. 2009;3(1):58-62.
18. Edak AU, Peter NC, Roseline CA, Victoria AA. Potentials of two Nigerian spices—*Piper nigrum* and *Monodora myristica* as sources for cheap natural antioxidants. American Journal of Plant Sciences. 2013; 4:1105-1115.
19. McCue PP, Shetty K. Inhibitory effects of rosmarinic acid extracts on porcine pancreatic amylase *in vitro*. Asian Pacific Journal of Clinical Nutrition. 2004;13:101-106.
20. Etouchi CB, Kuaté D, Ngondi JL, Oben J. Anti-amylase, anti-lipase and antioxidant effects of aqueous extracts of some Cameroonian spices. Journal of National Products. 2010;3:165-171.
21. Odo UE, Ezugwu CO, Ajali IU. Antimicrobial activity of *Monodora myristica* seed oil. Journal of Pharmacology and Allied Sciences. 2004; 2:233-236.
22. Ogu GI, Ekeanyanwu RC, Madagwu EC, Eboh OJ, Okoye J. *In vitro* antimicrobial evaluation of African nutmeg (*Monodora myristica*) seeds. International Journal of Tropical Agriculture and Food System. 2011;5:55-60
23. Ketosource: Exogenous ketones; 2017.
24. Iwu MM. Handbook of African medicinal plants. 1st Edn. CRC Press, Boca Raton, FL. 1993;205-2010.
25. Okwu DE. Phytochemicals, vitamins and mineral contents of two Nigerian medicinal plants. International Journal of Molecular Medicine and Advanced Science. 2005; 1(4):375-381.
26. Byju's the learning app; 2016.
27. Toshikazu K. Biological functions and health benefits of amino acids. Foods Food Ingredients Journal of Japan. 2002; 206.
28. The WHO Essential Medicines and Health Products Information; 2017.
29. Olga M, Lucia G, Michal Z, Tomas R, Jan M, Irena K. Using Odd-Alkanes as a carbon source to increase the content of nutritionally important fatty acids in *Candida krusei*, *Trichosporon cutaneum*, and *Yarrowia lipolytica*. Hindawi International Journal of Analytical Chemistry. 2017;9. Article ID: 8195329
Available:<https://doi.org/10.1155/2017/8195329>
30. "Ester." The Columbia Encyclopedia, 6th ed. *Encyclopedia.com*. 27 Mar. 2018.
Available:<http://www.encyclopedia.com>
31. Esters – Oils and Plants.
Available:www.oilsandplants.com
32. Gylling H, Plat J, Turley S, et al. Plant sterols and plant stanols in the management of dyslipidaemia and prevention of cardiovascular disease. Atherosclerosis. 2014;232(2):346–360.
33. Adewole E, Ajiboye BO, Idris OO, Ojo OA, Onikan A, Ogunmodede OT, et al. Phytochemical, Antimicrobial and Gc-Ms of African Nutmeg (*Monodora Myristica*). International Journal of Pharmaceutical Science Invention. 2013;2(5):25-32.
34. Enabulele SA, Oboh FOJ, Uwadiae EO. Antimicrobial, nutritional and phytochemical properties of *Monodora myristica* seeds. IOSR Journal of Pharmacy and Biological Sciences. 2014;9(4):01-06.

© 2018 Maduike and Anuna; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<http://www.sciedomain.org/review-history/24445>