

An Assessment of Hygiene Practices and Health Status of Street-food Vendors in Yaoundé, Cameroon

Nguendo Yongsy H. Blaise^{1*}

¹Health Geographer and Spatial Epidemiologist at Institute for Demographic Training and Research (IFORD), University of Yaoundé II, P.O.Box 1556 Yaoundé, Cameroon.

Author's contribution

This whole work was carried out by author NYHB.

Article Information

DOI: 10.9734/IJTDH/2014/10265

Editor(s):

(1) Anthony R. Mawson, Public Health and Institute of Epidemiology & Health Services Research, Jackson State University, USA.

Reviewers:

(1) Anonymous, University of N'Djamena, Chad.
(2) Claude Ahouangninou, Institut des Sciences Biomédicales Appliquées, Université d'Abomey Calavi, Cotonou, Bénin.

(3) Anonymous, Obafemi Awolowo University, Nigeria.

(4) Samir Mahgoub, Microbiology Department, Zagazig University, Egypt.

(5) Isaac Monney, Department of Environmental Health and Sanitation Education, University of Education, Winneba, Ghana.

Peer review History: <http://www.sciencedomain.org/review-history.php?iid=634&id=19&aid=5923>

Original Research Article

Received 21st March 2014
Accepted 24th May 2014
Published 1st September 2014

ABSTRACT

Background: From 1993, civil servants in Cameroon were subjected to a new working schedule, according to which, work starts every working day from 7.30am to 3.30pm with a short lunch break from 12 noon to 1.00pm. However, this new regulation was not accompanied by effective measures such as the creation of formal eating houses or canteens to ensure the provision of services that could satisfy the food needs of workers. Consequently, there has been a development and concentration of informal food vendors

*Corresponding author: Email: sir_nguendoyongsi@hotmail.com;

around public offices. While street foods are an important source of ready-to-eat nutritious, low-cost meals for the urban poor, the health risk posed by such foods may outweigh their benefits.

Objective: This study aims at assessing personal hygiene and health status of vendors involved in informal sales of food in Yaoundé.

Methods: In this cross-sectional epidemiological design, 837 food vendors were investigated through medical examinations notably stools analysis for ova and cysts of parasites. Questionnaires were also administered to the vendors. The vendors were selected from within the three main places in the city where people congregate: schools, market places, and government ministries/office.

Results: The personal hygiene of a majority of street-food vendors in Yaoundé (87, 1%) is of low and this is reflected on their health status because among the 837 food vendors enrolled in the study, 379 were found infected with one or more faeco-orally transmissible parasites, a 45% prevalence rate.

Conclusion: There is the need to improve the personal hygiene of food vendors in Cameroon in order to reduce the prevalence of food-related diseases. This can be done through adequate sensitization programmes on personal and food hygiene and the development of food handling policies.

Keywords: Food vendors; food consumption; informal sector; personal hygiene; oro-fecal transmitted diseases; parasites; health risk; Yaoundé.

1. INTRODUCTION

Urbanization in Africa has been phenomenal and puzzling; with a rapid shift from 15% in 1950 to about 41% urban proportion currently [1]. The UNFPA [2], estimates that by 2030, the continent may attain 54% urban proportion. This phenomenal growth has been qualified variously as “galloping” and “wild” [3] to express not only the uncontrolled nature, but also the socio-economic changes and sanitary consequences often associated with it [4]. Among those socio-economic changes, stands the upsurge of the “street food industry”. In fact, the street food sector is a source of nutrition, place of consumption and an important provider of income for many urban-based individuals and families in developing countries. It is widely recognized as an inevitable phenomenon tied to urban growth. In big cities, almost everyone (students, civil servants, traders, students and workers) eat street foods due not only to the long distances of workplaces from houses, but also because of their low cost. Street foods account for a major part of the daily diet and so contributes towards meeting nutritional requirements, although the contribution varies and is rarely quantified. Street foods are sold in almost every country in the world [5,6]. The FAO Alimentarius Commission defines “street foods” as ready-to-eat foods prepared and/or sold by vendors and hawkers especially in the streets and other public places [7]. Street foods show great variation in terms of ingredients, processing, methods of marketing and consumption. They often reflect traditional local cultures and exist in an endless variety, encompassing meals, drinks and snacks. There is much diversity in the raw materials as well as in the method of preparation of street foods. In addition, there are differences in the places where street foods are prepared. They can be broadly grouped into four categories namely: foods prepared in small-scale food factories or traditional workshops; foods prepared in the home; foods prepared in markets; and foods prepared on the street. While street foods are an important source of ready-to-eat nutritious, low-cost meals for the urban poor, the health risks posed by such foods may outweigh their benefits. In fact, certain street-vended foods pose significant risk to consumers due to microbiological contamination. Every year, millions of people become ill and thousands die

as a result of eating unsafe food [8]. The risks depend primarily on the type of food, the method of preparation and the manner in which it is held before consumption [9]. In addition to microbial hazards, several studies show that street foods can become contaminated with high levels of toxic chemicals including pesticide residues, heavy metals, *mycotoxins* and unapproved food additives [10,11]. Food can become contaminated at any point before its consumption, including during preparation. There is general acceptance on contamination and infection arising from practices of food preparation/storage/sale and consumer morbidity and even mortality. Explanations of health conditions of street vendors are still less debated, whereas there are many sicknesses which can be caused by inadequate/poor personal hygiene such as bacterial (*gastroenteritis*, diarrhea), viral (hepatitis A, colds and flu), and parasitic diseases (*giardiasis*, scabies infection, *pediculosis*, hookworm infection, and roundworm infection (*strongyloides*). Since there is a lack of awareness that poor personal hygiene practices can help the transmission of disease-causing germs [12], this work is intended to assess the personal hygiene of street-food vendors in Yaoundé as well as their health status. The paper focuses on disease-causing agents transmitted directly by the faeco-oral route.

2. MATERIALS AND METHODS

2.1 Definition of Concepts

2.1.1 Personal hygiene

First attested in English in 17th century, the word hygiene comes from the French *hygiene*, meaning "(art) of health" or "good for health" [13]. Thus, hygiene is an old concept related to medicine, as well as to personal and professional care practices related to most aspects of living. In medicine and in everyday life settings, hygiene practices are employed as preventative measures to reduce the incidence and spread of diseases. In line with this, the terms hygiene and cleanliness are often used interchangeably, which can cause confusion. In general, hygiene mostly means practices that prevent spread of disease-causing organisms [14]. Since cleaning processes such as hand washing and teeth brushing remove infectious microbes as well as dirt, they are often the means to achieve hygiene. Other uses of the term appear in phrases including: body hygiene, sleep hygiene, mental hygiene, dental hygiene, and occupational hygiene, used in connection with public health. Hygiene is also the name of a branch of science that deals with the promotion and preservation of health [15]. However, within the framework of this study, we are primarily interested in personal hygiene, which involves practices performed by an individual to care for one's body health and well being through cleanliness. Motivations for personal hygiene practice include reduction of personal illness, healing from personal illness, optimal health and sense of well being, social acceptance and prevention of spread of illnesses to others [16].

2.1.2 Health status

Health status is not a term that is commonly used or instantly understood. Most people, even those who are familiar with health care, consider it a medical jargon. The major reason that this term has eluded simple definition is the lack of an agreed-upon definition of health. An operational definition of health is essential before the health status of an individual or a population can be assessed. According to the WHO [17], health is "*the state of complete physical, mental and social well-being and not merely the absence of disease or infirmity*" and the "*extent to which an individual or group is able to realize aspirations and satisfy needs, and to change or cope with the environment. Health is a resource for everyday life,*

not the objective of living; it is a positive concept, emphasizing social and personal resources as well as physical capabilities." Although this definition has been subject to controversy as lacking operational value and because of the problem created by use of the word "complete," it remains the most enduring [18]. From an operational point of view, health is the level of functional or metabolic efficiency of a living organism. In humans, it is the general condition of a person's mind and body, usually meaning to be free from illness, injury or pain. Thus, since the main determinants of health include the social and economic environment, the physical environment, and the person's individual characteristics and behaviors [19], health status is the level of health, poor or good of the individual, group, or population. That level of health can be subjectively assessed by the individual or by more objective measures through indicators such as morbidity, mortality, and disability rates.

So explained, how then is health status measured in this study? Before presenting the method of measuring health status, it is important to conceptualize what is being measured. Health status needs to be examined in more detail in order to emphasize the complexity of health. (Fig. 1) below is a simple model of health status showing that health is the result of *health determinants* and of diverse interventions. Health status is defined through these two factors individually and as a result of the interactions between them. *Health determinants* laid down at the Ottawa conference in 1986 can be labeled proximal (direct) or distal (indirect) [20]. According to Lerer et al. [21], distal determinants mainly made up of "the macro-economic, educational, environmental, demographic and health factors, are considered to be the driving forces of health status and are often, but not invariably, mediated through proximal determinants". Besides, "proximal determinants whose some can be prevented (through health education/promotion) and regulated (through fiscal and legislative measures), are regarded as direct causes of changes in health status". Viewed as such, the relationship between determinants and health status is largely indirect and is modified only by factors like gender, education and socio-economic disparity.




Fig. 1. The relationship between health status, determinants and interventions

Thus, health is a multi-dimensional concept that is usually measured in terms of: (i) absence of physical pain, physical disability, or a condition that is likely to cause death, (ii) emotional well-being, and (iii) satisfactory social functioning. Some have advocated including the quality of an individual's physical environment in the definition of health, but this dimension is not at present included in the most widely used measures of health. However, there is no single "standard" measurement of health status for individuals or population groups. Individual health status may be measured by an observer, usually a physician who performs an examination and rates the individual along any of several dimensions, including presence or absence of life-threatening illness, risk factors for premature death, severity of disease,

and overall health. Individual health status may also be assessed by asking the person to report his/her health perceptions in the domains of interest, such as physical functioning, emotional well-being, pain or discomfort, and overall perception of health. Although it is theoretically fashionable to argue that the measurement of health should consist of the combination of both an objective component plus the individual's subjective impressions, no such measure has been developed. It is suggested that in the absence of comprehensive or absolute measures of the health of a population, average lifespan, prevalence of preventable diseases or deaths, and availability of health services serve as indicators of health status [22]. That is why most studies measure health status by using a variety of indices including morbidity, mortality, and available health resources. These health indicators are quantifiable characteristics of a population which researchers use as supporting evidences for describing the health status of a population. Thus, in the frame of this work, we have resorted to the morbidity indicator through the prevalence of parasites transmitted by the faeco-oral route.

2.2 Study Area

The study was conducted in Yaoundé, the capital city of Cameroon, situated in Central Africa between latitudes 3°47' and 3°56' North and 11°10' and 11°45' East (Fig. 2). Yaoundé displays the classical Equatorial climate characterized by regular and abundant rainfall of more than 1,600mm per annum, and a fairly high average annual temperature of 23°C. Like many sub-Saharan African cities, Yaoundé is currently experiencing very rapid urbanization. With an estimated annual growth rate of 4.5 per cent since 1980, its population has grown from 812,000 inhabitants in 1987 to 1,500,000 inhabitants in 2000, and to about 2,100,000 inhabitants in 2007. Thus, like many metropolises, Yaoundé is in the grip of sudden and unprecedented urban growth with an increase in the size of the city's labour force. Consequently, the demand for non-traditional services has increased and there has been a surge in service-oriented activities that are not part of the formal economic system. As the population pressure in the inner city grows, many people settle in suburbs and distant areas, and daily commuting has become a way of life. Civil servants and other formal sector workers who are subject to the 7.30am to 3.30pm work schedule (with short lunch break) have been forced to alter their eating schedules, attitudes and food tastes, by resorting to street-vended foods. Though foods sold on the streets are relatively cheap and readily available, lack of access to potable water supply and garbage disposal, and unsanitary environmental conditions further exacerbate the public health risk associated with street-food in such an urban setting.

2.3 Sources of Data and Methods of Collection

Data used in this work came from a cross-sectional study carried out on the town in 2013 by the working group Milieux, Environnements, et Développement Durable (MEDDU) of the Institute for Training and Research in Demography (IFORD). Although the whole city constituted the basis for sampling because of the proliferation of street food vending (even in gaps between buildings and in urban open spaces), a double strata sample was carried out. First, 3 main representative sectors were chosen out of the 11 where dwellers congregate daily: the City-center with its public offices and ministries, the Mokolo market which is the biggest one in Yaoundé, and the Ngoa Ekelle university campus with its faculties and colleges. Secondly, 837 food vendors were randomly selected from these 3 sectors on the basis that they sell food items that are either consumed directly or require cooking before consumption (Fig. 3). The survey was designed and supervised by the author but was carried out by nurses and professional investigators from IFORD. After four days of theoretical and practical training, those nurses and investigators visited targeted food

vendors to collect data through standardized questionnaires. The information collected related to age, educational qualification, knowledge, attitude and practices as regards parasite infestation), and to medical investigation (stools screening for identification of faeco-orally transmitted parasites). As far as ethical considerations are concerned, the study was first introduced to the National Ethics Committee which approved it giving the leeway for the data collectors to explain the purpose of the study to food vendors whose consent was obtained through a signed document for those who were willing to participate. Consented food vendors were given well labeled universal tubes in which they placed a little lump of freshly passed stool. Stool samples were later collected and transported to the parasitological unit of the Medical Laboratory Center located in the city center) within the accepted requirements, for (i) direct smear method and formol ether concentration methods as described by WHO [23], and clarified by Leventhal et al. [24]. (ii) and for confirmation and identification of parasite species. Computer programmes used were Epi info (for recording raw data, verifying and validating the data collected), and SPSS 10.0 for analysis and tabulation).




Fig. 2. Map of Cameroon showing the location of the urban community of Yaoundé

proper personal hygiene is of little consideration. (Table 3) shows practices of investigated food vendors with regard to their personal hygiene.

Table 1. Distribution of investigated food vendors in Yaoundé according to age and sex

Age group (years)	Food vendors				Total
	Males		Females		
	Frequency	%	Frequency	%	
[10–19]	57	16.38	49	10.03	106(12.66)
[20–29]	106	30.46	151	30.88	257(30.71)
[30–39]	130	37.36	195	39.87	325(38.84)
[40–49]	47	13.50	75	15.34	122(14.57)
[50–59]	8	02.30	19	03.88	27(03.22)
Total	348	100	489	100	837(100%)

Source: Field investigations, 2013

Table 2. Total revenues earned daily by food vendors in Yaoundé (in FCFA)

Revenue range	Areas of sale			Total
	Mokolo market	Around the ministries	University campus	
≤5 000	22(17.88%)	43(34.96%)	58(47.15%)	123(14.69%)
[5 000-10 000]	29(17.57%)	71(43.03%)	65(39.40%)	165(19.71%)
[10 000-5 000]	86(41.54%)	69(33.33%)	52(25.13%)	207(24.73%)
[15 000-20 000]	112(44.26%)	87(34.39%)	54(21.35%)	253(30.22%)
≥20 000	58(65.16%)	22(24.72%)	9(10.12%)	89(10.63%)

Source: Field investigations, 2013

The table above does not indicate who can be considered as foods vendors by virtue of personal hygiene. It just provides us with items that should help to build an index of personal hygiene quality. To build such an index, we must resort to the statistical technique called "score". It consists of classifying variables' modalities from "slightly poor personal hygiene" to "Very poor personal hygiene" (Table 4). Foods vendors with poor personal hygiene are those responding to at least four of the criteria and foods vendors with inadequate or very poor personal hygiene are those not responding for up to seven of the pre-defined criteria of adequacy which respond to more than six of the criteria. They represent 23.54% of our sample.

3.3 Assessment of Health Status of Street-food Vendors in Yaoundé

Among the 837 food vendors involved in the study and who were subjected to medical investigation, 379 were found infected with one or more faeco-orally transmissible parasites (i.e. a 45.28 percent prevalence rate). Men have appeared to be more vulnerable than women (Fig. 4) and independently of the gender, vendors less than 19 years were the most infected (Fig. 5).

Table 3. Personal hygiene practices of street-food vendors in Yaoundé

	Frequency	percentage
1. Bathing the body daily		
- No	309	38.10
- Yes	502	61.90
2. Washing hands regularly		
- No	683	81.61
- Yes	154	18.39
3. Brushing Teeth daily and regularly		
- No	196	23.91
- Yes	624	76.09
4. Wearing clean clothes daily		
- No	529	64.27
- Yes	294	35.73
5. Trimming or cutting Nails		
- No	612	77.87
- Yes	174	22.13
6. Hair cutting or shampooing		
- No	658	78.62
- Yes	179	21.38
7. Covering nose when coughing		
- No	741	93.37
- Yes	70	06.63
8. Wearing high/lace-up shoes		
- No	718	85.79
- Yes	119	14.21
9. Selling food in clean area?		
- No	385	46.00
- Yes	452	54.00

Source: Field investigations, 2013

Table 4. Building's process of the personal hygiene quality level

Table 4a. Hygiene quality (by sum of valid elements in increasing order)		
Valid	frequency	percentage
1	25	02.98
2	83	09.92
3	136	16.25
4	148	17.68
5	130	15.53
6	118	14.10
7	109	13.02
8	88	10.52
9	00	00.00
Total	837	100,00

Constitutive variables (nine): 1. teeth brushing, 2. nails cutting/trimming, 3. hand washing, 4. bathing, 5. wearing clean clothing, 6. wearing clean shoes, 7. covering nose when coughing, 8. hair cutting, 9. food handling in clean area. Average score:2,07; Median score:2,00

Table 4b . Personal hygiene according to level of quality or adequacy		
Level of quality	Distribution of the numbers	percentage
Slightly poor	108	12.90
Poor	532	63.56
Very poor	197	23.54
Coverage of the variable (837)	100.0	100.00

Source: Field investigations, 2013




Fig. 4. Distribution of Faecal orally transmissible parasites among street-food vendors in Yaoundé, according to gender




Fig. 5. Distribution of Faecal orally transmissible parasites among street-foods vendors in Yaoundé, according to age

Faeco-orally transmissible parasites found were the *Entamoeba*, and then followed the *Giardiasis*, *Ankylostoma duodenalis*, and *Ascaris lumbricoides* among others (Table 5).

Table 5. Frequency of selected parasite infections in street-food vendors in Yaoundé (n=379)

Species	Frequency	Percentage
<i>Entamoeba histolytica</i>	67	17.68
<i>Entamoeba coli</i>	61	16.10
<i>Giardia lamblia</i>	54	14.25
<i>Giardia duodenalis</i>	46	12.13
<i>Ankylostoma duodenalis</i>	49	12.93
<i>Ascaris lumbricoides</i>	43	11.34
<i>Trichuris trichiura</i>	42	11.08
<i>Enterobius vermicularis</i>	17	04.49
Coverage of the variable	100,0(379)	

Source: Microbiologic examinations operated at Laboratoire du Centre, Yaoundé, 2013

The study also investigated infections with multiple parasite species as a modulator of disease severity by examining the association of concomitant infection of parasites in the sample. And cases of polyparasitism were recorded in 248 vendors, i.e. 65.43%. There was more incidence of double infection compared to single or triple. Adults appeared to record more of single and double infection while teenagers recorded more double and triple infection (Table 6).

Table 6. Cases of polyparasitism among street-food vendors in Yaoundé, according to age group

Infection	Teenagers [10–19 years]	Young adults [20–39 years]	Adults [40 years and more]
Single	48(21.54%)	59(30.31%)	24(41.17%)
Double	91(50.77%)	75(48.48%)	21(52.94%)
Triple and more	39(27.69%)	19(21.21%)	3(05.89%)
Total	178(100%)	153(100%)	48(100%)

Source: Medical investigations, 2013

The results show that street food vendors with poor and inadequate (very poor) personal hygiene are more prone to faeco-oral transmissible parasites (35.09% and 46.70%) than those with slightly poor quality of personal hygiene (18.20%) (Table 7).

Table 7. Distribution of parasites species according to quality level of street-food vendors personal hygiene in Yaoundé

Parasite species	Quality level of personal hygiene		
	Slightly poor	Poor	Very poor
<i>Entamoeba histolytica</i> (n=67)	11(15.94%)	23(17.29%)	33(18.65%)
<i>Entamoeba coli</i> (n=61)	13(18.85%)	19(14.28%)	29(16.38%)
<i>Giardia lamblia</i> (n=54)	9(13.05%)	20(15.03%)	25(14.12%)
<i>Giardia duodenalis</i> (n=46)	12(17.39%)	19(14.28%)	15(08.47%)
<i>Ankylostoma duodenalis</i> (n=49)	7(10.14%)	14(10.52%)	28(15.81%)
<i>Ascaris lumbricoides</i> (n=43)	7(10.15%)	18(13.55%)	18(10.18%)
<i>Trichuris trichiura</i> (n=42)	8(11.59%)	14(10.52%)	20(11.30%)
<i>Enterobius vermicularis</i> (n=17)	2(02.89%)	6(04.53%)	9(05.09%)
Total	69(100.00%)	133(100.00%)	177(100.00%)

Level of significance: $p < 0.005$ Source: Field and medical investigations, 2013

Not surprisingly, results obtained show that high exposure to faeco-oral transmissible parasites are informal settlements, particularly the peri-urban and Sub-central spontaneous quarters areas known as to be poorly equipped with socio-economic infrastructure (Table 8).

Table 8. Spatial variation of personal hygiene quality of street-food vendors in Yaoundé according to type of their quarter

Type of quarters	Level of quality of personal hygiene						Total
	Slightly poor		Poor		Very poor		
	Frequency	%	Frequency	%	Frequency	%	
Wealthy residential quarters	00	00	00	00	00	00	00
Housing estates	13	12.04	27	05.07	03	01.52	43 (05.13)
Communal plots	17	15.74	59	11.10	14	07.11	90 (10.75)
central spontaneous quarters	25	23.14	83	15.60	46	23.35	154 (18.41)
Sub-central spontaneous quarters	32	29.63	138	25.94	75	38.07	245 (29.27)
Peri urban quarters	15	13.89	130	24.44	40	20.31	185 (22.10)
Semi rural quarters	6	05.56	95	17.85	19	09.64	120 (14.34)
Total	108	100	532	100	197	100	837(100%)

Level of significance: $p < 0.002$; Source: Field and medical investigations, 2013

4. DISCUSSION

The street food sector is very important in Cameroonian cities for socio-cultural and economic reasons: deteriorating living conditions in rural areas, rapid urbanization, structural adjustment and economic recession [25]. However, the activity accounts for 20 to 30 percent of household expenditure, and is also a major provider of employment. The street food sector has therefore received special attention worldwide, when its socio-cultural and sanitary relevance was finally recognized. Street foods have often been seen as the sources of the outbreak of cholera in most developing countries. This perception triggered priority sanitary actions to reinforce street food control programmes among which was the evaluation of vendors' personal hygiene. Yet, what is considered proper personal hygiene can be culture-specific and may change over time. In some cultures, the removal of body hair is considered proper hygiene. In others, kissing or shaking of hands to reduce transmission of bacteria by contact is hygienic. Other personal hygienic practices would include covering one's mouth when coughing, disposal of soiled tissues appropriately, making sure toilets are clean, and making sure food handling areas are clean, besides other practices [26]. This means that personal hygiene practices should be assessed according to specific criteria like those used in Bangkok [27], Calcutta [28], in Lima [29], and in Johannesburg [30]. In this study, food vendors with adequate personal hygiene would have been those responding to our nine criteria, that is, who bath daily in the morning, brush their teeth regularly, wash their hands with soap all the times, cut their nails and hair, wear clean clothes and high shoes, and cover their noses and mouths with a tissue or the hand when coughing or sneezing, and place food on clean platforms. Unfortunately, we didn't find food

vendors who fulfilled all those criteria. Besides, we found foods vendors who respond to seven of our nine criteria of adequacy and therefore are hereby considered as vendors with slightly adequate personal hygiene. However, health issues related to street foods are related to disease transmission route. In fact, the human body provides places for disease-causing germs and parasites to grow and multiply. These places include the skin and in and around the openings to the body. Parasites are very common around the world and infections are transmitted in crowded places. Intestinal parasites are spread when faecal matter gets into the mouth. This can happen through ingesting contaminated food or water. Hookworms like *ancylostoma* are generally contracted by walking barefoot on soil contaminated by faeces from infected persons or by wading in contaminated water. In a nutshell, poor sanitation and unsafe water increase the risk of catching parasites. It is less likely that germs and parasites will get inside the body if people have good personal hygiene habits. Using the guide to good personal hygiene drafted by the Australian government [31], personal hygiene was measured in this study through the following practices: washing the body, washing hands regularly and especially before handling food, cleaning the teeth at least once a day, wearing clean clothing, covering the nose and mouth with a tissue or the hand when coughing or sneezing, making sure food handling areas are clean, cutting or trimming finger nails, keeping hair short or removing hair, wearing lace-up/high shoes. We did not include items like availability of toilet facilities, access to water system closets, use of tap water, waste disposal considered by Costa-Cruz and Cardoso [32] and Muinde and Kuria [33] because they pertain to environmental hygiene or to community health [34,35]. Unsafe food has been a human health problem and many food safety problems encountered today are not new. Although governments throughout the world are doing their best to improve safety of food supply, occurrence of food borne disease remains a significant health issue in both developed and developing countries. While the food borne disease burden is often most well-documented in developed countries, the implications for developing countries are extremely serious in terms of both human suffering and increased demand on health care systems. The long term consequences are malnutrition and increased vulnerability to a wide range of diseases. This is particularly true for faecal orally transmissible diseases. The prevalence of faecal-orally transmissible parasite recorded in this study (45.28%) is indicative of a high level of faecal contamination of food since these parasites are acquired via accidental consumption of parasite eggs and cysts [36]. This prevalence signals high possibility of transmission of these parasites through infected food vendors to their customers especially in cases where such food vendors are unhygienic in food handling. The facts that those food handlers are asymptomatic make them cysts and/or egg passers and are unaware that they are possible transmitters of parasitic infections/diseases. Yet, they are and are accounted to the WHO evidence that, every year, millions of people become ill and thousands die as a result of eating unsafe food. These foods have come to be contaminated at any point before its consumption, mostly during preparation and delivering by vendors whose personal hygiene has been assessed to be insufficient or poor.

A large proportion of street-food vendors in Yaoundé has attained the secondary education i.e. have acquired basic education on hygienic practices, but like it has been reported by Ghosh et al. [op cit] in India, they lack adequate means and an enabling environment to safely serve the clients. The three types of quarters/suburbs with higher percentages of infected food vendors were central spontaneous quarters, sub-central spontaneous quarters and peri urban quarters with poor living conditions (overcrowded rooms, lack of adequate sanitation, low or limited connection to the drinking water network). This may be due to the lack of adequate supply of water which could encourage food vendors to bath at least in the morning before going out for vending, if we agree with Kubota et al. [37] that bathing in the

morning is important from a hygiene point of view as it removes the bacteria that form by the process of sweating and that which settle down from external sources, and therefore appears as a preventative measure to reduce the incidence and spread of disease. In those suburbs, majority of the food vendors use the pit latrine and other related structures and/or dung hills for defaecation. These structures may however, facilitate contact with stools from which they can either acquire infection or carry eggs and cysts of parasites with which contamination of their wares may be inevitable thereby leading to widespread contamination of food and drinks. In addition, lack of public toilets in their points of sale may be one of the factors of epidemiology since food hawkers make use of dung hill for defaecation while hawking their wares. The exposure of food items by stationery food vendors facilitated contact of insect with such food items, thereby increasing possible contamination of foods with eggs and cyst of parasites since insects have been identified as carriers' of eggs and cysts of parasites [38]. Furthermore, those overpopulated quarters contain the major slums in the city. And due to high cost of living and high rental cost, most vendors live in shack houses that are situated in marshy areas with increased probability of acquiring faecal-oral infections [39]. From our field observations, one of the most critical problems in street food vendors in Yaoundé is their indecent physical appearances: uncut hairs and nails, sandals or dirty plastic shoes, clothes soaked with sweat. This is a sign of their low standards of living which prevent them from taking care of their bodies. According to the FAO and Akindes [40,41], the ultimate objective of the street vendors is not to take care of their body, but to earn their living.

This study shows the high incidence of protozoan parasites (nearly 60%) among food vendors and a high prevalence of *Entamoeba histolytica* and *Entamoeba coli*. Poverty may be responsible for this prevalence and/or re-infection with these parasites since the use of soap in washing of hands for example, is considered an additional cost even by the adult vendors that are aware of the need for hygienic practices in food handling. From the results, we note that none of the food vendors met the WHO requirements for effective hand washing which include washing in soapy water before preparing or handling foods. The lower incidence of multiple infections among adult food vendors implies that the routine inspection and supposed periodic de-worming have a reducing effect on infection especially with helminthes [42]. Children and teenagers are known to be more vulnerable to parasitic infection than adults because of their level of exposure to parasite eggs and cysts, their level of hygiene and immunity to infection [43,44]. This may account for higher prevalence of infection recorded among teenagers' food vendors. Infection with helminthes recorded indicate high level of environmental contamination with helminth eggs, coupled with the fact that faecal-oral mode of transmission is the easiest and commonest mode of transmission of parasites in tropical areas [45]. In such a situation, the use of anti-helminthes drugs is less effective if factors promoting transmission have not been removed.

5. CONCLUSION

Street foods are ready-to-eat foods prepared and/or sold by vendors and hawkers especially in the streets and other public places. The street food sector plays an important role in providing accessible, low-cost meals for urban populations, particularly those in many developing countries. In Cameroon, they often reflect traditional local cultures and exist in an endless variety encompassing meals, drinks and snacks. The street food sector also plays an important role in providing employment opportunities for millions of men and women with limited education or skills, especially as the initial investment is low. In Yaoundé as well as in Douala, the main Cameroonian metropolises, street vending has even become an important component of food distribution system, particularly for midday meals. However, certain

street-food vendors can pose significant risk to consumers due to microbiological contamination. In fact, vendors are often poorly educated and untrained in food safety. Yet, they work under unsanitary conditions with little or no infrastructure support, but their personal hygiene is generally inadequate. Research has shown that their health status is also bad, since some of them are suffering from faecal-oral transmitted parasites or infections. This study reveals the need to improve the personal hygiene of food vendors in Cameroon in order to reduce the prevalence of faecal-oral transmitted parasites. Based on this, there is the need for adequate enlightenment programmes for street food handlers. There is also the need to enact and implement appropriate food handling policies. While some countries have conducted training programmes to educate street vendors on food safety, the development of training materials has to be tailored to meet their needs and situations. Recently the Five Keys of Safer Food have been successfully utilized in an evidence-based training programme for the vendors to improve their food handling practices. In 2009, WHO published a Train the trainer guide on the Five Keys to Safer Food which builds upon the safe food handling behaviors developed for the Five Keys to Safer Food Poster. To our knowledge, this study through its cross sectional design and direct investigation is the first to examine the extent to which faecal-oral transmitted parasites may be mediators in the relationship between personal hygiene and health outcomes in Cameroon. However, there are two main limitations to the study. First, the cross sectional design does not allow us to infer that any associations was causal. Second, we only modeled the relationships between quality of personal hygiene and faecal-oral transmitted parasites and intermediate health outcome. While we were underpowered to address more distal outcomes, faecal-oral transmitted parasites are closely linked to unsafe drinking water, and contaminated foods. If our results were to be replicated with other populations in different settings and across health conditions and outcomes, the use of a multi-level design with all known pathogens of faecal-oral transmitted pathogens would be recommended.

CONSENT

Not applicable.

ETHICAL APPROVAL

Not applicable.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. United Nations. World urbanization prospects: The 2005 revision. Washington, Department of economic and social affairs, population division. 2005;196.
2. United Nations Population Fund (UNFPA). State of world population 2007- Unleashing the potential of urban growth; 2007.
3. Sietchiping R. Geographic information systems and cellular automata-based model of informal settlement growth. PhD Dissertation, School of Anthropology, Geography and Environmental Studies, University of Melbourne. 2003;276.

4. Trang DT, Molbak K, Phung DC, Dalsgaard A. Helminth infections among people using wastewater and human excreta in periurban agriculture and aquaculture in Hanoi, Vietnam. *Tropical Medicine & International Health*. 2007;12:82–90.
5. Chakravarty, Canet C. Street foods in Calcutta. *Food, Nutrition and Agriculture*. Roma, FAO. 1996;17/18:23.
6. Hutabarat LSR. Street foods in Bangkok: The nutritional contribution and the contaminants content of street foods. An Andre Mayer research fellowship study. FAO, Rome. 1994;30.
7. FAO. Street foods. Report of an FAO Expert Consultation, Jogjakarta, Indonesia. FAO Food Nutr. Pap. 1989;46:1-96.
8. WHO. Prevention of food-borne disease: The five keys to safer food. Geneva, International food safety authorities network. 2006 ;4.
9. OMS. Essential safety of foods sold on the street. Unpublished manuscript. WHO / FNU / FOS 96.7 Revised Edition. Geneva.
Available: http://whqlibdoc.who.int/hq/1996/WHO_FNU_FOS_96.7_Rev1_fre.pdf.
10. WHO. Chemical risks in food. Geneva, Initiative to estimate the Global Burden of Foodborne Diseases. Retrieved from: <http://www.who.int/foodsafety/chem/en/>. [Consulted on April 2014]
11. FAO/OMS. Draft code of hygienic practice for the preparation and sale of street foods. Programme mixte FAO/OMS sur les normes alimentaires. Alinorm 91/15, Appendix III. Rome; 1990.
12. Ghosh M, et al. Prevalence of enterotoxigenic *Staphylococcus aureus* and *Shigella* spp. in some raw street vended Indian food. *Int. J. Environ. Health Res*. 2007;17:151-6.
13. Liddell HG, Scott R, Stuart HJ, Mckenzie RA. Greek-English Lexicon: Good for the health, wholesome, sound, healthy. Revised and augmented throughout by. Sir Henry Stuart Jones with the assistance of Roderick McKenzie. Oxford. Clarendon Press; 1940.
14. Australian government. Personal hygiene for food handlers. accessed from the Queensland Health. Retrieved from: www.health.qld.gov.au/foodsafety. Consulted on April 2014.
15. Aiello AE, Coulborn RM, Perez V, Larson EL. Effect of hand hygiene on infectious disease risk in the community setting: a meta-analysis. *American Journal of Public Health*. 2008;98:1372-81.
16. Lindsay Dingwall Personal Hygiene Care. John Wiley & Sons. 2010;336.
17. WHO. Constitution of the World Health Organization–Basic Documents, Forty-fifth edition, Supplement, October; 2006.
18. Jadad AR, O'grady L. How should health be defined? *British Medical Journal*. 2008;337:43-55
19. WHO. The determinants of health. Geneva. Retrieved from: www.who.int, and consulted on march; 2014.
20. Mikkonen J, Raphael D. Social Determinant of Health: The Canadian Facts. Toronto, York University, School of Health policy and management. 2010;35.
21. Lerer LB, et al. Health for all: Analyzing health status and determinants. *World Health statistics Quaterly*. 1998;51(1):7-20.
22. Bergner M. Health status measures: An overview and guide for selection. *Am. Rev. Publ. Health*, 1987;8:191-208.
23. WHO. Basic laboratory methods in medical parasitology. Geneva. 1981;69.
24. Leventhal R, Russell F, Cheadle. *Medical Parasitology: A Self-Instructional Text*. F.A. Davis Company, 6th edition. 2012;272.

25. Nguendo yongsi HB, Ntetu Lutumba A. Household standard of living and children's health in Sub-Saharan cities. *Journal of Developing Areas*. 2008;42(1):251-259.
26. Bloomfield SF, Stanwell-Smith R, Crevel RWR, Pickup J. Too clean, or not too clean: The Hygiene Hypothesis and home hygiene. *Clinical and Experimental Allergy*. 2006;36:402-25.
27. Hutabarat LSR. Street foods in Bangkok-the nutritional contribution and the contaminants content of street foods. Roma, FAO/An André Mayer Research Fellowship Study; 1994.
28. Chakravarty, Canet C. Street foods in Calcutta. *Food, Nutrition and Agriculture*. Roma, FAO. 1996;17/18:23.
29. FAO. Low-cost urban food distribution systems in Latin America. Roma, selected paper number. 2008;111:57.
30. Van Tonder I, Lues Jan FR., Theron, Maria M. The personal and general hygiene practices of food handlers in the delicatessen sections of retail outlets in South Africa. *Journal of Environmental Health*, 2007, pp 38-54
31. Australian Government/department of health. Environmental health program management and community education. In *Environmental Health Practitioner Manual: A resource manual for Environmental Health Practitioners working with Aboriginal and Torres Strait Islander Communities*. Available:<http://www.health.gov.au/internet/publications/publishing.nsf/Content/ohp-enhealth-manual-atsi-cnt-l~ohp-enhealth-manual-atsi-cnt-l-ch7>. Cnsulted on january 2014.
32. Costa-Cruz JM, Cardoso ML. Intestinal parasites in cooks and their helps in the city of Uberlandia, Minas Gerais Brazil. *Rev Institute Medical Tropical Soa Paulo*. 1997;38:100-107.
33. Muinde OK, Kuria E. Hygiene and sanitary practices of vendors of street foods in Nairobi, Kenya, *African Journal Of Food Agriculture Nutrition And Development*. 2005;7:1-15.
34. Hémond D. Epidemiological research on environment and health: Some methodological aspects. *Journal of Epidemiology and Public Health*. 1995;43(5):395-411.
35. Gérin M, Gosselin P, Cordier S, Viau C, Quenel P, Dewailly É. *Environment and Public Health-Principles and Practices*, Paris, Editions Tec & Doc. 2003;1022.
36. Ghosh M, et al. Prevalence of enterotoxigenic *Staphylococcus aureus* and *Shigella* spp. in some raw street vended Indian food. *Int. J. Environ. Health Res*. 2007;17(2):151-156.
37. Kubota K, Tamura K, Take H, Kurabayashi H, Mori M, Shirakura T. Dependence on very hot hot-spring bathing in a refractory case of atopic dermatitis. *Journal of medicine*. 1994;5:333-336.
38. Umeche N. Parasite ova and cysts in fruits in Calabar, Nigeria. *The Nigeria Journal of Parasit*. 1991;12:85-87.
39. Aiello AE, Coulborn RM, Perez V, Larson EL. Effect of hand hygiene on infectious disease risk in the community setting: a meta-analysis. *American Journal of Public health*. 2008;98:1372-1381.
40. FAO/Food Basket Foundation International. Study on street foods in Nigeria: comparative study of the socio-economic characteristics of food vendors and consumers in Ibadan, Lagos and Kaduna. Ibadan; 1991.
41. Akindes F. Urbanization and development of informal food sector in Côte d'Ivoire: The example of Abidjan. Paris, EHESS, Orstom. 1990;357.
42. Borkow G, Bentwich Z. Eradication of helminthic infections may be essential for successful vaccination against HIV and tuberculosis. *Bulletin of the World Health Organization*. 2000;78(11).

43. Montresor, et al. Helminth control in school-age children: A guide for managers of control programs. World Health Organization; 2002.
44. Luby S, Agboatalla M, Feikin DR, Painter J, Billhimmer W, Atref A, Hoekstra RM. Effect of hand-washing on child health: A randomized control trial. *Lancet*. 2005;366:225–233.
45. Nguendo Yongsi HB, Tejiokem CM, Tsague DL. Care activities in a semi-urban clinic in Cameroon (Nkondjock): Review of thirty years of parasitological examinations and prescription. *Black African Medicine*. 2007;54(1):79-90.

© 2014 Blaise; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history.php?iid=634&id=19&aid=5923>