



## **Emergence of Garlic (*Allium sativum* L.) as Influenced by Low Storage Temperature and Gibberellic Acid Treatments**

**Fikrte Woldeyes<sup>1\*</sup>, Kebede W/tsadik<sup>2</sup> and Getachew Tabor<sup>3</sup>**

<sup>1</sup>Department of Plant Science, Collage of Agriculture, Madda Walabu University, P.O.Box 247, Bale Robe, Ethiopia.

<sup>2</sup>Department of Plant Science, Collage of Agriculture and Environmental Science, Haramaya University, P.O.Box 138, Haramaya, Ethiopia.

<sup>3</sup>Agricultural Research Center, Debrezeit, Ethiopia.

### **Authors' contributions**

This work was carried out in collaboration between all authors. Author FW designed, analyzed, interpreted and prepared the manuscript. Authors KW and GT corrected the manuscript. All authors read and approved the final manuscript.

### **Article Information**

DOI: 10.9734/JAERI/2017/29843

Editor(s):

(1)

(2)

Reviewers:

(1)

(2)

(3)

Complete Peer review History:

**Original Research Article**

**Received 30<sup>th</sup> September 2016**

**Accepted 24<sup>th</sup> November 2016**

**Published 23<sup>rd</sup> December 2016**

### **ABSTRACT**

The cost of garlic cloves together with inconsistent emergence due to dormancy issues which needs rest or dormant period of more than three months, has contributed to the reduction of garlic production cycle in a year. Thus, pre-planting garlic clove treatment with Gibberellic acid (GA<sub>3</sub>) plant hormone and cold temperature storage duration are an important agronomic concern for dormancy breaking of local garlic (*Allium sativum* L.) variety. Greenhouse experiment was conducted in the spring cropping season of 2013 to investigate the effect of GA<sub>3</sub> concentration (0, 125, 250 and 375 ppm) and cold storage (7°C) durations (10, 20, and 30 days) on emergence of local garlic cultivar. The treatments were laid out in factorial arrangement with three replications using completely randomized design. The interaction effect of GA<sub>3</sub> and cold storage duration

\*Corresponding author: E-mail: fikr2119@gmail.com;

significantly increased on emergence percentage (96.66%), emergence rate (1.09) and number of sprouted cloves 0.447(100%) at GA<sub>3</sub> 125 ppm for 20 days of cold storage duration. In conclusion the result of the experiment revealed that treatment GA<sub>3</sub> 125 ppm with 20 days of cold storage duration is required for attaining optimum emergence performance of the local garlic variety in Haramaya Eastern Ethiopia.

*Keywords: Cold temperature; gibberellic acid and emergence of garlic.*

## 1. INTRODUCTION

Garlic (*Allium sativum* L.) is a member of *Alliaceae* family. It is an erect biennial herb normally grown as annual crop. It originated in the northwestern side of the Tien-Shan Mountains of Kirgizia in the arid and semi-arid areas of central Asia [1]. It is used for seasoning in many foods worldwide. It also has higher nutritive value compared to other bulb crops in addition to antibiotics like garlicin and allistatin it contains [2].

In tropical Africa, garlic is grown during the cold season in the Sahel region and high elevations in east Africa and in South Africa [3]. In Ethiopia, the total area under garlic production in 2012 was 21,558 ha and production was about 222,548 metric ton with average yield of 10.5t/ha. Garlic is produced for home consumption and as a source of income to many peasant farmers in many parts of the country [4]. The per capita consumption of garlic crops was estimated at about 1.74 kg and 5.9 kg in the rural and urban areas, respectively [5].

Even though garlic production throughout the year under rain-fed and irrigation conditions is possible, its potential has not been fully exploited due to scarcity of planting material, in addition to genetic, environmental and management factors. Garlic production is limited by planting material scarcity, due to the seed clove requiring resting period of more than three months after harvest as well as relatively short shelf life that do not sustain cloves to next cropping season. Garlic cloves require more than three months of ambient temperature and relative humidity storage for more uniform sprouting [6]. Despite beneficial effects on storing and availability of bulbs for human consumption, garlic dormancy creates problems for immediate use as planting material [7]. Use of cloves as planting material before completion of their innate dormancy period leads to garlic uneven sprouting.

Clove sprouting and emergence are controlled mainly by temperature and its storage period

[8,9]. During storage, sucrose and starch are consumed at high content and a part of non-consumed carbohydrates causes glucose accumulation of garlic tissue, enhancing sprouting [10]. Cloves treated at 10°C for 30 days storage temperature increased sprouting and percentage of emergence [11]. The inductive effect of cold storage depends not just on temperature regime but also on its duration depending on cold requirements for breaking bulb dormancy [12].

Plant growth regulators have been also known to play a vital role in sprouting of garlic [13]. Gibberellic acid is today known as a very efficient stimulating factor for breaking dormancy and increasing sprouting in many agronomic and horticultural crops [14].

The promoting effect of GA<sub>3</sub> treatment is often attributed to mobilization of stored reserves [15]. It was observed that the percentages of sprouting were higher at 250 ppm than at 125 or 500 ppm [13].

Farmers normally use garlic cloves as planting material after long storage period at room temperature that extend its dormancy to the next cropping year. Thus, it restricts the production cycle twice a year under rain fed and irrigation condition. Future prospects of Ethiopian small scale and large scale irrigation scheme is to increase vegetable production, including garlic, through increasing the frequency of production cycle per year. Thus, an increase in demand for garlic planting material is expected. To supplement this using freshly harvested garlic clove seed as a planting material after treating with growth promoter hormones and low storage temperature would be imperative.

Many studies have indicated that the application of growth regulators and low temperature storage period can shorten dormancy period of bulb crops, including garlic. However, there is scarce information regarding the effects of Gibberellic acid and low storage temperature treatments on sprouting of local garlic cultivars. Thus, the

objective of this study was to evaluate the effectiveness of application of different concentrations of GA<sub>3</sub> and low storage temperature on emergence of garlic.

## 2. MATERIALS AND METHODS

### 2.1 Experimental Site Description

The research was conducted at Haramaya University glasshouse research site during 2013 spring cropping season. The study area is located in the Eastern Hararghe zone of Oromia Regional State, 14 km west of Harar town and 508 km east of Addis Ababa, Ethiopia. Geographically, it is located between latitudes 9° 24' 53.13" N and 9° 24' 51.34" N, and longitudes 42° 01'55.69" E and 42° 01' 56.62" E at an altitude of 1980 meter above sea level.

### 2.2 Treatments and Experimental Design

The experiment consisted of 4 x 3 factorial combinations of Gibberellic acid concentrations (0, 125, 250 and 375 ppm) and storage durations (10, 20, and 30 days) stored at low temperature (7°C) (Table 1). The experiment was carried out in a glasshouse to evaluate sprouting and emergence parameters of garlic seed cloves. The treatments were laid out as a completely randomized design in a factorial arrangement where each treatment was replicated three times.

### 2.3 Experimental Procedures

Uniform and fresh bulbs of a local garlic variety, harvested at physiological maturity, were obtained from a farm in Haramaya University area. The bulbs were cured for one week and

separated into cloves. The cloves were stored at low temperature storage (7°C) for 10, 20, and 30 days. Then those stored cloves were soaked in Gibberellic acid solutions at concentrations of 0,125, 250, and 375 ppm for 24 hours. Gibberellic acid solution was made by dissolving Gibberellic acid powder in a drop of 70% ethanol enough to wet the powder for each concentration. Then it was mixed with distill water to get the required treatment concentrations.

A controlled experiment was undertaken in a glasshouse. A total of 20 cloves were planted on a seed tray per experimental unit. Planting was made on bench filled with seedbed media (3:2:1 soil, manure and sand mix ratio, respectively). Then proper watering of the seedbed was done until the trial was terminated. Sprouted and emerged cloves were recorded daily starting from one week after planting until one and half month, the period when cloves were expected to completely emerged.

### 2.4 Data Collection

Data on sprouted and unsprouted cloves and days to emergence were recorded from all planted cloves (20 cloves) in each experimental unit. The cloves establishment count was determined as percentages and rate of emerged cloves. As suggested by [16], emergence percentage (%) and emergence rate was recorded for each treatment as follows:

#### 2.4.1 Emergence (%)

Determined as (Number of normal seedlings per total number of cloves planted) \*100.

**Table 1. Treatments used in the study**

No	Treatments		
	Gibberellic acid (ppm)	Cold temperature storage (Days)	Combinations
1	0	10	GA <sub>3</sub> 0 x Day 10
2	0	20	GA <sub>3</sub> 0 x Day 20
3	0	30	GA <sub>3</sub> 0 x Day 30
4	125	10	GA <sub>3</sub> 125 x Day 10
5	125	20	GA <sub>3</sub> 125 x Day 20
6	125	30	GA <sub>3</sub> 125 x Day 30
7	250	10	GA <sub>3</sub> 250 x Day 10
8	250	20	GA <sub>3</sub> 250 x Day 20
9	250	30	GA <sub>3</sub> 250 x Day 30
10	375	10	GA <sub>3</sub> 375 x Day 10
11	375	20	GA <sub>3</sub> 375 x Day 20
12	375	30	GA <sub>3</sub> 375 x Day 30

### **2.4.2 Emergence rate**

Determined as Number of normal seedlings/ (days of first count) +...+ Number of normal seedlings / (days of final count).

### **2.4.3 Sprouted cloves**

It was determined as the sum total of all sprouted cloves.

### **2.4.4 Unsprouted cloves**

It was determined as the sum total of all non-sprouted cloves (Total number planted – total number sprouted).

## **2.5 Statistical Analysis**

Data collected for all parameters were subjected to analysis of variance (ANOVA) using statistical analysis system institute package (SAS, 2010 version 9.1). Comparisons of treatment mean values were made by least significant difference (Fisher's LSD) and Duncan's Multiple Range Test (DMRT) at 5% probability.

## **3. RESULTS AND DISCUSSION**

### **3.1 Sprouted Cloves**

Data presented in Table 2 indicated that the interaction effects of GA<sub>3</sub> and cold storage duration was highly significant (P< 0.01) for the number of sprouted cloves of garlic. The interaction effect data showed that combinations of GA<sub>3</sub> at 125 ppm and 20 days of cold storage, GA<sub>3</sub> at 250 ppm with 20 and 30 days of cold storage were equally effective in enhancing sprouting of all cloves of garlic planted, 0.447 (100%). Cloves that received 250 GA<sub>3</sub> for 10 days cold storage treatments gave the lowest percentage, 0.420 (88.33%) of sprouted cloves. In general, cloves treated with GA<sub>3</sub> at 250 ppm, except those treated with cold for 10 days, appeared to be better in improving percent sprouted cloves. This result is in agreement with those of [13] who observed that the percentages of sprouting in garlic were higher at 250 ppm than at 125 ppm or 500 ppm. Similarly, author [14] found that Gibberellic acid is a very efficient stimulating factor for breaking dormancy and increasing sprouting in many horticultural crops.

Cold storage of cloves for 20 or 30 days also significantly increased sprouting, especially with supplement of GA<sub>3</sub> at 250 ppm. Temperature below 10 was effective in breaking dormancy of

garlic [17]. Similarly, [11] reported that at storage of 10°C for 30 days increased sprouting of cloves. Garlic sprouting at low temperature is attributed to a complex of biochemical reactions which is carried out by interposition of different enzymes [18,19] that increase glucose content associated with sprouting, as glucose increment means sprouting initiation of garlic [10,18].

### **3.2 Unsprouted Cloves**

The percentage of unsprouted cloves 0.141(10.00%) was found to be higher in the treatment 10 days of cold treatment alone, 125 ppm GA<sub>3</sub> concentrations with 30 days prior cold storage treatments and 375 ppm GA<sub>3</sub> concentrations with 10 days of cold storage treatment. This might be due to dormancy in freshly harvested garlic bulbs whose sprouting depends on air temperature and storage period [18,19] as well ineffectiveness of low rates of GA<sub>3</sub> to replace cold treatment or probably toxicity at increased dose of GA<sub>3</sub> [20,21].

### **3.3 Emergence Percentage (%)**

The results obtained from the greenhouse experiment exhibited that the effect of GA<sub>3</sub> concentration and its interaction with cold storage duration were highly significant (P<0.01) and the effect of cold storage duration was also significant (P<0.05) on seedling emergence percentage of garlic (Table 2).

Cloves stored under cold for 20 days and treated with GA<sub>3</sub> 125 ppm gave the highest percentage (96.66%) of seedling emergence, which was statistically similar to values recorded in all cold stored cloves for 20 or 30 days regardless of GA<sub>3</sub> treatment, except those stored for 30 days and treated with either 125 ppm or 375 ppm GA<sub>3</sub>. Moreover, combination GA<sub>3</sub> 125 and 250 ppm with 10 days of cold storage had significantly similar percentage of germination. On the other hand, cloves treated with GA<sub>3</sub> 125 ppm with 30 days of cold storage had the lowest (66.66%) seedling emergence percentage.

In general, cloves treated with GA<sub>3</sub> 250 ppm at all level of cold storage duration showed better seedling emergence percentage. This result agreed with observation of [13] who reported that concentrations of GA<sub>3</sub> in the same magnitude significantly improved percentage emergence in garlic. From the effect of cold storage duration extending from 10 to 30 days improved emergence percentage of garlic seedlings that agree with results of [11] who reported that the

**Table 2. Interaction effects of Gibberellic acid and cold storage temperature duration on seedling emergence percentage (%) and rate, sprouted and unsprouted cloves of garlic**

Treatment	Sprouted cloves (%)		Un sprouted cloves (%)		Emergence	
	Transformed Mean	Original Mean	Transformed Mean	Original Mean	Percentage	Rate
GA <sub>3</sub> 0 x Day 10	0.435c	95.00c	0.100e	5.00e	71.66cd	0.53def
GA <sub>3</sub> 0 x Day 20	0.424de	90.00de	0.141cd	10.00a	83.33a-d	0.61de
GA <sub>3</sub> 0 x Day 30	0.443ab	98.33ab	0.033f	1.66f	91.66ab	0.86bc
GA <sub>3</sub> 125 x Day 10	0.439bc	96.66bc	0.100e	5.00e	85.00abc	0.56def
GA <sub>3</sub> 125 x Day 20	0.447a	100.00a	0.000g	0.00f	96.66a	1.09a
GA <sub>3</sub> 125 x Day 30	0.424de	90.00de	0.141cd	10.00a	66.66de	0.42efg
GA <sub>3</sub> 250 x Day 10	0.420ef	88.33ef	0.152cd	11.66ab	85.00abc	0.69cd
GA <sub>3</sub> 250 x Day 20	0.447a	100.00a	0.000g	0.00f	88.33abc	0.56def
GA <sub>3</sub> 250 x Day 30	0.447a	100.00a	0.000g	0.00f	80.00a-d	0.84bc
GA <sub>3</sub> 375 x Day 10	0.424de	90.00de	0.141cd	10.00a	78.33bcd	0.61de
GA <sub>3</sub> 375 x Day 20	0.435c	95.00c	0.100e	5.00e	80.00a-d	0.51d-g
GA <sub>3</sub> 375 x Day 30	0.428d	91.66d	0.127de	8.33cd	76.66bcd	0.45efg
LSD(0.05)	0.0067**	3.17**	0.0276**	2.94**	17.14 **	0.20 **
CV (%)	0.9	2.04	18.0	25.71	13.03	19.31

\*\* significantly different at 1% probability level; Means followed by the same letter within a column are not significantly different, at 5 % or 1% levels of significance (LSD); CV=coefficient of variation. Data transformed by square root

highest percentage of emergence was found in bulbs stored at 10°C for 30 days. Similar with this study, [22] reported that pre-chilled cloves at 4°C for three weeks gave increased emergence percentage (95.6%) than non-chilled (74.6%) cloves of garlic. Cold storage for 30 and 40 days reduced the dormant period and increased field emergence [23]. The proportion of seed clove germination was raised by the use of gibberellins and/or cold treatment could probably be due to rise of amino acid content in embryo, which is the source hydrolytic enzyme needed for digestion of endospermic starch when seeds restart growth at germination [24,15,25]. Such greater seedling emergence by GA<sub>3</sub> and/or cold storage treatments would enable faster crop development, early maturity and higher yields.

### 3.4 Emergence Rate

Significant difference on emergence rate of garlic seed clove were observed due to GA<sub>3</sub> concentration (P<0.01), cold storage duration (P<0.05) and their interaction (p<0.01) (Table 2). Cold storage of cloves for 20 days and treatment with GA<sub>3</sub> at 125 ppm resulted on the greatest seedling emergence rate (1.09). The lowest seedling emergence rate (0.42) was recorded from cloves that treated for 30 days cold storage with GA<sub>3</sub> at 125 ppm treatment. Increase in cold storage durations also increased at the rate of seedling emergence when no GA<sub>3</sub> was applied. The higher value indicated that 30 days cold storage duration had good ability to break garlic

dormancy earlier which could be important for successful field establishment as well as good yield than late emerged cloves. In agreement with the present observation [9,18] reported that seedling emergence in garlic has been enhanced by low temperature treatments. Storage of garlic at temperatures of 5 to 10°C than 25°C had promoted respiration rate during storage and the emergence of seedlings [10].

This exhibited that GA<sub>3</sub> could replace the cold storage requirement of dormant garlic for early sprouting and other early growth of the crop.

The result corroborates earlier reports by [24,15,25] that indicated presence of adequate levels of this acid, which accumulates in dormant seeds during cold storage and/or when exogenously applied, stimulates the synthesis, activation and secretion of hydrolytic enzymes, releasing reducing sugars and amino acids which are essential for embryo growth.

### 4. CONCLUSIONS

Greenhouse experiment was conducted to determine the effect of cold storage duration and GA<sub>3</sub> concentration on emergence performance of garlic. The results showed that the combination of 125 ppm GA<sub>3</sub> with cold storage for 20 days resulted in a highly significant difference in all of those emergence parameters. GA<sub>3</sub> at 250 ppm combined with 20 and 30 days of cold storage produced highly significant improved number of

sprouted cloves while extending cold storage duration alone highly significantly improved emergence percentage and rate.

Generally, the results of the present study showed highly significant effects of GA<sub>3</sub> concentration and cold storage duration on emergence of garlic. It could, thus, be concluded that using treatments of GA<sub>3</sub> at 125 ppm with cold storage duration at 20 days could be optimum for breaking dormancy of the local garlic variety and production of two cropping a year under rain-fed and irrigation conditions around Haremaya for the aim of supplementing planting material. However, it is difficult to reach a conclusive recommendation since the experiment was conducted at one location and one season. Therefore, this research should be repeated at different area using improved and high-yielding local varieties of garlic with clove cutting and other dormancy breaking methods is suggested for possibility of improving emergence of fresh seed cloves.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Etoh T, Simon WP. Diversity, fertility and seed production in garlic. In: Rabinowitch HD, Currah L, editors. *Allium crop science: Recent advances*. Wallingford: CABI Publishing; 2002.
2. Mahmood N. Horticultural crops production; 2000. Available: [www.cabi.org/about-cabi/staff/P0033](http://www.cabi.org/about-cabi/staff/P0033) (Accessed 24 May 2014)
3. Grubben GH. Plant resources of tropical Africa vegetables. *Plant Resources of Tropical Africa*. 2004;2:56–61.
4. Getachew T, Asfaw Z. Achievements in Shallot and Garlic Research Report. Ethiopian Agricultural Research Organization; 2000.
5. Metasebia M, Shimelis H. Proceedings of the 15<sup>th</sup> annual research and extension review meeting, Alemaya University of agriculture, Ethiopia; 1998.
6. Getachew T, Getahun. Asfaw Z. Influence of storage duration on field sprouting, maturity and yield of some garlic (*Allium sativum* L.) cultivars at Debre Zeit, Ethiopia. *Journal of Horticultural Science and Biotechnology*. 2004;79 (6):871-876.
7. Ledesma A, Reale MI, Racca R, Burba JL. Effect of low temperatures and pre planting storage time on garlic clonal type Rosado Paraguayo growth. *Phyton (Argentina)*. 1980;39:37-48.
8. Takagi H. Garlic *Allium sativum* L. In: Brewster JL, Rabinowitch HD. editors. *Onions and allied crops*. Florida: CRC Press, Ltd; 1990.
9. Del Pozo A, Gonzalez MI, Barraza C. Phenological development of thirty clones of garlic (*Allium sativum* L.) influence of temperature, photoperiod and cold storage. *Acta Horticulturae*. 1997;433: 389–394.
10. Contwell MI, Kang J, Hong G. Heat treatments control sprouting and rooting of garlic cloves. *Postharvest Biological Technology*. 2003;30:57-65.
11. Youssef SN. Growth and bulbing of garlic as influenced by low temperature and storage period treatments. *Journal of Rural Observation*. 2013;5(2):47-56.
12. Del Pozo A, Gonzalez G. Developmental responses of garlic to temperature and photoperiod. *Agricultural Technology*. 2005; 65:119-126.
13. Rahman HM, Haquel SM, Karim AM, Ahmed A. Effects of Gibberellic acid on breaking dormancy in garlic (*Allium sativum* L.). *International Journal of Agriculture and Biology*. 2008;1560(1):63–65.
14. Faghihi G, Radan M. Side effects of herbal drugs used in dermatologic disorders. *Journal of Cosmetics, Dermatological Sciences and Applications*. 2011;1(1):1-3.
15. Soyler D, Khawar KM. Seed germination of caper (*Capparis ovate* var. *herbacea*) using  $\alpha$  naphthalene acetic acid and Gibberellic acid. *International Journal of Agricultural Biology*. 2007;9:35–37.
16. Maguire JD. Rate of germination-aid in selection and evaluation for seedling emergence and vigor. *Crop Science*. 1962; 2:176-177.
17. Rahman HM, Muhammad SH, Masood A. Pre-planting temperature treatments for breaking dormancy of garlic cloves. *Asian Journal of Plant Science*. 2003;2:123-126.
18. Vazquez BM, Lopez EG, Mercado SE, Castano TE, Leon VG, Ratindo KE. Low temperature storage of garlic for spring planting. *Horticultural Science*. 2006; 39(3):571.
19. Atashi SV, Akbarpor K, Mashayekh, Mousavizadeh SJ. Garlic physiological

- characteristics from harvest to sprouting in response to low temperature. *Journal of Stored Products and Postharvest Research*. 2011;2(15):285–291.
20. Masud AM. Effects of planting date, growth regulators and cold treatments on the growth and yield of garlic. Master of Science Thesis, Bangladesh Agricultural University, Mymensingh, Bangladesh; 1995.
  21. Rahman SM, Islam AM, Haque SM, Karim AM. Effects of planting date and Gibberellic acid on the growth and yield of garlic (*Allium sativum L.*). *Asian Journal of Plant Science*. 2004;3(3):344-352.
  22. Moustafa YM. Performance of new imported foreign garlic genotypes grown under the Egyptian conditions. *Egypt Journal of Agriculture Research*. 2011; 87(1):219-243.
  23. Silva NF, Casali VWD. Effects of low temperature storage and planting date on dormancy in the garlic cultivar *Peruano*. *Horticultural Abstracts*. 1987;59:320.
  24. Mayer AM, Poljakoff MA. *The Germination of Seeds*. 4<sup>th</sup> ed. UK: Pergamon Press; 1989.
  25. Tayyaba H, Khalid H, Saba S, Ahtisham U. Effect of Gibberellic acid (GA<sub>3</sub>) on morphological and physiological attributes of ispaghol (*Plantago ovata L.*). *International Journal of Water Resources and Arid Environments*. 2012;2(6): 190-196.

© 2017 Woldeyes et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://sciencedomain.org/review-history/17324>